

Towards laser based improved experimental schemes for multiphoton $e^+ e^-$ pair production from vacuum.

I. Ploumistakis[†], S. D. Moustazis[†], and I. Tsohantjis[‡],

Technical University of Crete, Department of Sciences,

[†]*Institute of Matter Structure and Laser Physics,*

[‡]*Division of Mathematics*

Chania GR-73100, Crete, Greece

(Dated: July 15, 2009)

Numerical estimates for pair production from vacuum in the presence of strong electromagnetic fields are derived, for two experimental schemes : the first concerns a laser based X-FEL and the other imitates the E144 experiment. The approximation adopted in this work is that of two level multiphoton on resonance. Utilizing achievable values of laser beam parameters, an enhanced production efficiency of up to 10^{11} and 10^{15} pairs can be obtained, for the two schemes respectively.

PACS numbers: 12.20.-m, 42.50.Hz, 42.55.-f

Keywords: Pair production, Multiphoton processes, High intensity lasers

I. INTRODUCTION

Electron-positron pair production from vacuum breakdown in the presence of strong electromagnetic fields is one of the most interesting non-linear QED phenomena and at the same time a rich area of theoretical and experimental investigations [1, 2, 3, 4, 5, 6, 7, 8]. Recently the possibility for an experimental verification of the phenomenon is plausible due to the rapid development of ultra-intense laser facilities [11, 12, 13]. The first steps on the theoretical treatment of the phenomenon can be found [1], but Schwinger [2] was the first that examined it thoroughly. By the implementation of the proper time method Schwinger obtained the following conditions for pair creation to occur : The invariant quantities $\mathcal{F} = \frac{1}{4}F_{\mu\nu}F^{\mu\nu} = -\frac{1}{2}(\vec{\mathcal{E}}^2 - c^2\vec{\mathcal{B}}^2)$, $\mathcal{G} = \frac{1}{4}F_{\mu\nu}\tilde{F}^{\mu\nu} = c\vec{\mathcal{E}} \cdot \vec{\mathcal{B}}$, where $F_{\mu\nu}$ and $\tilde{F}_{\mu\nu} = \frac{1}{2}\epsilon_{\mu\nu\alpha\beta}F^{\alpha\beta}$ are the electromagnetic field tensor and its dual respectively, must be such that neither $\mathcal{F} = 0$, $\mathcal{G} = 0$ nor $\mathcal{F} > 0$, $\mathcal{G} = 0$. These restrictions are for example satisfied close to the antinodes of a standing wave or at the region of a focused laser beam. In order to have sizable effects the electric field strength must exceed the critical value $\mathcal{E}_c = \frac{mc^2}{e\lambda_c} \simeq 1.3 \times 10^{18} \text{V/m}$ which corresponds to focal laser intensities of the order of 10^{29}W/cm^2 and above. Such laser intensities can be reached in the very near future from facilities such as [14], [29], [30] and XFEL [18]. Brezin and Itzykson [3] and soon after Popov [6] used multiphoton atom ionization technics such as the imaginary time method [6, 9, 10] showed that the probability of pair creation over a 4- Compton volume can be obtained as

$$w_P = \sum_{n>n_0} w_n \quad (1)$$

With the parameter of the theoretical treatments being $\gamma = mc\omega/e\mathcal{E} = \hbar\omega\mathcal{E}_c/mc^2\mathcal{E}$, which is the equivalent of Keldysh parameter, the two important areas $\gamma \ll 1$ (high electric field strength and low frequency, where the adiabatic non-perturbative tunneling mechanism dominates) and $\gamma \gg 1$ (low electric field strength and high frequency, multiphoton mechanism), lead respectively to the following number of pairs $N(\tau)$ [6] :

$$N(\tau) = 2^{-3/2}n_0^4(\mathcal{E}/\mathcal{E}_c)^{\frac{5}{2}}\exp\left(-\frac{\pi\mathcal{E}_c}{\mathcal{E}}\left(1 - \frac{1}{2\left(n_0\frac{\mathcal{E}}{\mathcal{E}_c}\right)}\right)\right)(\omega\tau/2\pi), \quad \gamma \ll 1$$

$$N(\tau) \approx 2\pi n_0^{3/2}\left(\frac{4\gamma}{e}\right)^{-2n_0}(\omega\tau/2\pi), \quad \gamma \gg 1$$

with τ being the pulse duration and $n_0 = 2mc^2/\hbar\omega$. The first experimental verification of e^-e^+ pair production took place at SLAC (E-144 experiment)[18]. Highly energetic electrons (maximum 46.6GeV) underwent nonlinear Compton scattering with n laser photons (of $\lambda = 527 \text{nm}$, $\omega = 2.35 \text{eV}$ and laser pulses of energy 500mJ), $e + n\omega \rightarrow e' + \omega'$, producing backscattered photons ω' of $27 - 30 \text{GeV}$ which then collide with the laser photons to produce e^- , e^+ pairs, via the multiphoton Breit -Wheeler mechanism, $\omega' + n\omega \rightarrow e^- e^+$. The number of positrons measured in 21962 laser pulses was 175 ± 13 for a $n = 5.1 \pm 0.2$ (statistical) $_{-0.8}^{+0.5}$ (systematic) multiphoton order process, a result that is in very good agreement with the theory. These results have been shown in [20] to be obtained also from Popov's theory.

In a recent paper Avetissian et al [19] treated e^-e^+ pair production in a standing wave $\mathbf{A} = 2\mathbf{A}_0 \cos \omega t$ formed by oppositely directed laser beams of plane transverse linearly polarized electromagnetic waves of frequency ω , using a two level on resonance multiphoton approximation [19, 20, 21]. The main difference between this approach and the previous one is the condition of resonance, which as shown in the detailed investigation presented in [21, 22, 23, 24], can lead to very high production efficiency. The probability density of pair creation for the case of oscillating electric field, is given in relation (33) in [19]. This approximation will be considered in the rest of the paper. Following [19, 21], taking the momentum of the created electrons(positrons) to be $\mathbf{p} = (p_x = p \sin \theta, p_y = p \cos \theta, 0)$, where θ is angle between $\mathbf{A}_0$ (lying on the Oy axis) and $\mathbf{p}$, the probability density w_{n0} and the number of pairs created N_0 at angle $\theta = 0$ (where both maximize) for an n th order process are respectively given by [21, 22, 23]

$$w_{n0} = \frac{n\omega}{8\pi^2} f_{n0}^2 (n^2\omega^2 - 4m^2)^{\frac{1}{2}} \quad (2)$$

$$N_0 = w_{n0} V \tau = \frac{1}{4\pi^2} \frac{V\tau}{V_e} \frac{q\sqrt{q^2 - 1}}{m^2 c^4} f_{n0}^2 \quad (3)$$

In deriving (2) and (3) the resonance condition

$$n = 2E/\hbar\omega = 2qmc^2/\hbar\omega, \quad q \geq 1 \quad (4)$$

has been used and

$$f_{n0} = \frac{E}{4p_y} \left(1 - \frac{p^2}{E^2}\right)^{\frac{1}{2}} n\hbar\omega J_n\left(4\xi \frac{mp}{E\omega}\right).$$

$\xi \sim \frac{1}{\gamma}$ is the relativistic invariant parameter given by

$$\xi = \frac{mc^2 \mathcal{E}_0}{\hbar\omega \mathcal{E}_c}, \quad (5)$$

$\mathcal{E}_0$ is the amplitude of one incident wave and $\xi \lesssim 1$ for the approximation to hold. $V_e = 7.4 \times 10^{-59} m^3 s$ is the 4-Compton volume of the electron, $V \sim \sigma^2 l$ is the interaction volume, τ is the pulse duration, σ is the cross section radius and $l \ll \lambda$ the electromagnetic wavelength [19, 21]. Finally since in the Bessel function $J_n(x)$, $x \simeq n$, the approximation $J_n(n \operatorname{sech} \alpha) = (1/\sqrt{2\pi n \tanh \alpha}) (a = \operatorname{sech}^{-1}(\frac{2\xi}{q} (1 - \frac{1}{q^2})^{\frac{1}{2}}))$ has to be used giving,

$$f_{n0} = \frac{1}{4} (q^2 - 1)^{-\frac{1}{2}} n\hbar\omega \frac{\exp(n \tanh a - na)}{\sqrt{2\pi n \tanh a}} \quad (6)$$

As shown in [21], relation (3) leads to very good estimates for the theory both when higher harmonics are implemented and when the range of its applicability is examined. For example for a Nd-Yag laser with $\omega = 1.17eV$, $\lambda = 1\mu m$, intensity $1.35 \times 10^{22} W/m^2$, $\tau \sim 10^{-14} s$, $\sigma \sim 10^{-5} m$, $l = 0.1\lambda$, the fifth harmonic can create approximately up to 10^{12} pairs for $n \sim 10^5$, provided that the electric field strength does not exceed the order of $10^{13} V/m$.

In section two that follows we present the use of a table top laser based X-ray Free Electron Laser scheme for pair creation and in section three a scheme analogous to the one used in the E-144 experiment. The aim of this paper is to emphasize the advantages of using such schemes for the experimental observation of the on resonance approximation [19, 21], as they can be realized in well equipped laboratory and can provide satisfactory results which we will show on our numerical estimations that will follow. One should mention that the following two setups allow the increase of electric field strength $\mathcal{E}$ with a simultaneous increase of ω , so that ξ remains $\lesssim 1$, as it is essential for the validity of the multiphoton on resonance approximation.

II. PAIR CREATION USING LASER BASED X -FEL SYSTEM

In this section we investigate the use of what one could call a table top laser based X-ray Free Electron Laser for electron positron pair creation. Such system is similar to a standard X -FEL system [17] with the main difference between the two to be the electron beam creation and acceleration mechanism. The table top XFEL utilizes a

relativistic electron beam produced and accelerated by a high intensity ($\sim 10^{20} - 10^{21} W/cm^2$) ultrashort laser beam [25, 26, 27, 28]. Then the electron beam propagates through a wiggler system and X-ray beam is created. Generation of such relativistic electron beam from a laser source can be achieved by interaction with a solid or gas target and is a wide field of research. For our purposes we consider a laser system with intensity of $10^{21} W/cm^2$ and pulse duration $20 fs - 30 fs$. Recent experiments [25, 26, 27, 28] have confirmed that the accelerated electron beam can reach energies E_{beam} close to $1 GeV$ and charge close to $1 nCb$, but is expected that in the near future this charge will become available. This beam enters an undulator with period λ_u and after the interaction with the magnetic field in it, X-ray photons of wavelength

$$\lambda = \frac{\lambda_u}{2\gamma_L^2} \quad (7)$$

will be produced, where γ_L is the relativistic factor ($\gamma_L = \frac{E_{beam}}{m_e c^2}$). The undulator period for our scheme is $\lambda_u = 5 mm$ but it can be adjusted depending on the X-ray photon wavelength we wish to produce.

Using (7) for $E_{beam} = 1 GeV$ we obtain X-ray beam wavelength $\lambda = 0.65 nm$, which corresponds to photon energy $\omega = 1.909 KeV$. Two opposite directed X-ray beams form a standing wave to a circular spot area of radius $\sigma = 100 nm$. The interaction 4-volume in which pair creation takes place is taken as $V = \sigma^2(0.1\lambda)\tau$, where $\tau = 100 fs$. We also assume that the conversion efficiency of the electron beam energy to X-ray photons is about 10% and given the power of the electron beam $P_{beam} = 1 \times 10^{13} W$, we obtain the energy of the X-ray beam $E_b = 0.1 J$. As in [21], with the above parameter values, we firstly investigate, using (3) the dependence of pair number N_0 to q (electron rest mass units). The result is presented in Figure 1, where the envelope of $N_o(q)$ is plotted, for $\xi = 0.9532$. This optimum choice of ξ (leading to high production efficiency), is dictated by energy conservation between the X-ray beam energy E_b and the energy $E = 2qmc^2 N_0$ of created pairs: one can numerically solve the energy conservation equation $E_b = 0.1 J = 2qmc^2 N_0(\xi, q)$ for appropriate range of values of the q parameter (see [21] for a detailed analysis on the choice of ξ).




FIG. 1: Envelope of created $e^+ e^-$ pair number N_o vs rest mass unit q for $\xi = 0.9632$ and photon energy $1.909 keV$.

The most probable pair number is the one that corresponds to $q = 1.40$ and is of the order of 10^{10} pairs. The multiphoton order of the process as obtained by (4) is $n = 750$.

Secondly one can also investigate the dependence of N_o from ξ (equivalently $\mathcal{E}_0$). Using (3) and the data from Table 1 we present in Figure 2, the dependence of pair number N_0 from ξ for three E_{beam} values of $1 GeV$, $400 MeV$ and $200 MeV$, the last two being considered in order to estimate and compare pair creation efficiencies for energies that can be easily created with the recent technology in large laser facilities. Note that in order to obtain each of the three plots in Figure 2 the values of (q, n) are respectively $(1.41, 750)$, $(1.40, 4742)$, $(1.40, 18833)$.

Each one of the curves has a different ending point, because of previous mentioned energy conservation consideration. The choice of plotting vs. parameter ξ is justifiable, since it depends both from the electric field strength $\mathcal{E}_0$ and photon energy and thus allowing to investigate the dependence of N_0 on both of these physical quantities. By adjusting the photon energy and/or the electric field one can estimate the number of produced pairs for the corresponding value of ξ from figures like Figure 2. Observe the very sensitivity of N_0 from ξ which is a result of resonance approximation (this is due to the presence of the Bessel function in rel (3)).




FIG. 2: Log- plot of the number of created pairs N_0 vs ξ for electron beam energy $1GeV$ (top curve), $400MeV$ (middle curve) and $200MeV$ (bottom)

Electron beam energy E_{beam}	Wavelength of X ray laser beam (nm)	X-ray photon energy (eV)	Maximum Electric field Strength (V/m)
$1GeV$	0.65	1909	3.4×10^{15}
$400MeV$	4.06	305	5.5×10^{14}
$200MeV$	16.3	76	1.4×10^{14}

Table 1. Typical parameters for obtaining Figure 2

The electron beam charges considered for these energies case are respectively 1nC for 1GeV , 0.4 nC for 400MeV and 0.2 nC for 200MeV. The estimates of Figure 2 show that even if we choose to lower the electron beam energy, we can achieve efficient electron positron pair production (of the order of 10^{11} for 1GeV , 10^{10} for 400 and 200 MeV) and strongly suggest the possible use of this scheme for experimental observation of resonance approximation.

III. PAIR CREATION USING A CONFIGURATION ANALOGOUS TO THE E144 EXPERIMENT.

The second scheme proposed is based on the E144 experiment at SLAC [18].It consists of two steps. On the first step a relativistic electron beam is produced and accelerated by an ultrashort laser beam of photon energy 1eV. On the second step the relativistic electron beam interacts with the main high intensity ($10^{20} W/cm^2$) highly focused laser beam. On the electron's frame the electric field intensity of the laser beam is thus increased due to the relativistic factor while the focusing guarantees that the condition $\mathcal{E}^2 - H^2 > 0$, for pair creation is satisfied. On the electrons reference frame the electric field $\mathcal{E}^*$ and the photon energy $E_{photons}^*$ are given by

$$\mathcal{E}^* = \gamma_L \mathcal{E}_{laserlab} \quad (8)$$

$$E_{photons}^* = \gamma_L E_{photons} \quad (9)$$

where $\gamma_L = \frac{E_{beam}}{m_e c^2}$, E_{beam} being the electron beam energy.

For $E_{beam} = 1GeV$, we obtain from (9) $E_{photons}^* = 1.956 keV$, which corresponds to $\lambda = 0.1nm$ and $\mathcal{E}^* \sim 10^{15}V/m$. We will apply the same analysis as in the section two , to estimate the efficiency of resonance approximation for this scheme. Using (3) where the interaction 4-Volume $V = \sigma^2(0.1\lambda)\tau$, $\lambda = 0.1nm$, $\sigma = 50\mu m$ and $\tau = 10^{-14}s$, we present in Figure 3 the envelope of the created pair number N_0 as a function of q for $\xi = 0.9632$ (ξ is obtained with same considerations as before).

The most probable process corresponds to $q = 1.41$ and can produce up to the order of 10^{16} pairs. The multiphoton order of this process is $n = 735$. We also examine the efficiency of pair creation for two more energies of the electron beam, $400MeV$ and $200MeV$. As in section two, using formula (3) and the data from Table 2 we present in Figure 4, the dependence of pair number N_0 from ξ for three E_{beam} values of $1GeV$, $400MeV$ and $200MeV$ with corresponding values (q, n) for the later two energies as $(1.41, 1844)$ and $(1.40, 3686)$




FIG. 3: Envelope of created $e^+ e^-$ pair number N_o vs rest mass unit q for $\xi = 0.9632$ and photon energy 1.956 keV .




FIG. 4: Log- plot of the number of created pairs N_0 vs ξ for electron beam energy 1 GeV (top curve), 400 MeV (middle curve) and 200 MeV (bottom)

Electron beam energy E_{e-beam}	X-ray photon energy (eV)	Electric field Strength (V/m)
1 GeV	1909	4.7×10^{15}
400 MeV	782	7.8×10^{14}
200 MeV	391	1.9×10^{14}

Table 2. Typical parameters for obtaining Figure 4

The estimates of Figure 4 of this scheme strongly suggest the possible use of this scheme for experimental observation of resonance approximation (of the order of 10^{16} for 1 GeV , 10^{15} for 400 and 200 MeV).

IV. CONCLUSION

The above analysis aims to support the experimental verification of e^+e^- pair creation using the on resonance multiphoton approximation by presenting two possible schemes. One is based on using a table-top X-FEL laser system and the second is analogous to the E144 experiment. The numerical estimates presented that a sufficient number of pairs can be produced by both schemes that can reach up to the order of 10^{11} pairs for the table top X-FEL and 10^{15} for the E144 like scheme. An important fact in both proposed schemes is that they use laser based modern technology for production and acceleration of electron beam and consequently they do not need any large acceleration facilities.

Moreover, since the existing laser laboratories have this kind of technology, the required experimental values of the parameters needed for pair production can be achieved. Our choice to carry out our numerical estimates with E_{beam} no greater than 1 GeV is thus justified, as electron beams with energies like 400 MeV or 200 MeV can be created quite easily. Besides the above advantages of the two schemes, the laser beam parameters can be easily adjusted to change the electron beam energy, in order to operate on resonance. Moreover the criterion for the choices of the values of the parameters used, is that they should be close to typical values rather than extreme ones, in order to maintain our estimates as realistic as possible. Consequently our analysis, suggests that both of the schemes presented are promising for a future experimental verification of the on resonance pair production, especially when new powerful laser systems like X-FEL[17], HIPER[29] and ELI[30] are under development.

-
- [1] O. Klein, Z. Phys., **53**, 157 (1929); F. Sauter, Z. Phys. **69**, 742 (1931); W. Heisenberg, H. Euler, Z. Phys. **98**, 718 (1936).
 - [2] J. W. Schwinger, Phys. Rev., **82**, 664 (1951).
 - [3] E. Brezin and C. Itzykson, Phys. Rev. D **2**, 1191 (1970).
 - [4] G.J. Troup and H.S. Perlman, Phys. Rev. D **6**, 2299 (1972).
 - [5] W. Greiner, B. Muller, J. Rafelski, 'Quantum Electrodynamics of Strong Fields', Springer – Verlag, Berlin, 1985.
 - [6] V.S. Popov, JETP Lett. **13**, 185 (1971); Sov. Phys. JETP **34**, 709 (1972); Sov. Phys. JETP **35**, 659 (1972); V.S. Popov and M. S. Marinov, Sov. J. Nucl. Phys. **16**, 449 (1973) ; JETP Lett. **18**, 255 (1974); Sov. J. Nucl. Phys., **19**, 584 (1974).
 - [7] A. I. Nikishov, Nucl. Phys. B **21**, 346 (1970).
 - [8] N.B. Narozhnyi and A. I. Nikishov, Sov. J. Nucl. Phys. **11**, 596 (1970); Sov. Phys. JETP, **38**, 427 (1974).
 - [9] A. Ringwald, Phys. Let. B **510**, 107 (2001).
 - [10] V.S. Popov, Phys. Let. A **298**, 83 (2002).
 - [11] M. Perry and G. Mourou, Science **264**, 917 (1994)
 - [12] P. Chen and C. Pellegrini, in Quantum Aspects of Beam Physics, Proc.15th Advanced ICFA Beam Dynamics Workshop, Monterey, Cal., 4-9 Jan 1998 (World Scientific, Singapore, 1998) p. 571.
 - [13] P. Chen and T. Tajima, Phys. Rev. Lett. **83**, 256 (1999).
 - [14] T. Tajima and G. Mourou, Phys. Rev. STAccel.Beams **5**, 031301 (2002)
 - [15] S. S. Bulanov, N.B. Narozhny, V.D. Mur, and V.S. Popov, Phys. Lett. A **330**, 1 (2004)
 - [16] D. B. Blaschke, A.V. Prozorkevich, C. D. Roberts, S. M. Schmidt and S. A. Smolyansky, Phys. Rev. Lett, **96**, 140402 (2006).
 - [17] XFEL-Technical Design Report (2006) Publisher: DESY XFEL Project Group, European XFEL Project Team, Deutsches Elektronen-Synchrotron, Member of the Helmholtz Association, Notkestrasse 85, 22607 Hamburg, Germany ISBN 3-935702-17-5
 - [18] D.L. Burke et. al., Phys. Rev. Let., **79**, 1626 (1997).
 - [19] H. K. Avetissian, A. K. Avetissian, G. F. Mkrtchian and Kh. V. Sedrakian, Phys. Rev. E **66**, 016502 (2002).
 - [20] C. Kaberidis, I. Tsohantjis and S. Moustazis in 'Frontiers of Fundamental and Computational Physics' Udine, Italy, 26-29 September 2004, Sidharth B.G, Honsell F., de Angelis A. (Eds.) 2005 pp. 279-283
 - [21] I. Tsohantjis, S. Moustazis and I. Ploumistakis, Physics Letters B **650** 249(2007)
 - [22] I. Ploumistakis, Master Thesis Technical University of Crete, (2007)
 - [23] I. Ploumistakis, I. Tsohantjis and S. Moustazis, "New approaches on Laser Vacuum Breakdown for Pair Creation", 35th EPS Conference on Plasma Phys. Hersonissos, 9 - 13 June 2008 ECA Vol.32D, P-1.123 (2008)
 - [24] I.Tsohantjis, S. D. Moustazis, I. Ploumistakis, "Pair creation from vacuum in the presence of ultra-intense laser beams", 35th EPS Conference on Plasma Phys. Hersonissos, 9 - 13 June 2008 ECA Vol.32D, O-4.041 (2008)
 - [25] V. Malka, J. Faure, Y.Glinec, A. Pukhov, J.Rousseau Phys. of Plasmas **12**, 056702 (2005)
 - [26] F. Amiranoff et.al Phys. Rev. Lett. **81**, 995 (1998)
 - [27] W.P Leemans, et.al Nature Physics **2**, (2006)
 - [28] S.P.D. Mangles, et.al Nature **431**, 535 (2004)
 - [29] <http://www.hiper-laser.org>
 - [30] <http://www.extreme-light-infrastructure.eu>